



Nobivac[®] Myxo-RHD PLUS

SWITCHING RABBITS THAT HAVE BEEN PREVIOUSLY VACCINATED ONLY WITH NOBIVAC MYXO-RHD (INCLUDING THOSE THAT HAVE LAPSED (>12 MONTHS) WITH VACCINATION FOR AN INACTIVATED RHDV-2 VACCINE)

Nobivac Myxo-RHD PLUS will be replacing Nobivac Myxo-RHD, which will no longer be available once all current stock has been exhausted.

Nobivac Myxo-RHD PLUS has proven efficacy in naïve rabbits and effectively boosts rabbits previously vaccinated against myxomatosis, RHDV-1 and RHDV-2. However, please note that Nobivac Myxo-RHD PLUS vaccine product labelling also includes the following statement:

Rabbits that have been vaccinated previously with another myxomatosis vaccine, or that have experienced natural myxomatosis infection in the field, may not develop an adequate immune response against rabbit haemorrhagic disease following vaccination.

This statement relates to the phenomenon of vector interference which is very relevant to a very specific subset of rabbits being switched over to Nobivac Myxo-RHD PLUS. Based on the results of efficacy and switching studies, an optimum approach to avoid this potential risk in all relevant groups is laid out overleaf.

Nobivac[®] 
Protection unites us.

Nobivac[®] Myxo-RHD PLUS

GROUP 1

Naïve rabbits



Vaccinate with Nobivac Myxo-RHD PLUS from 5 weeks of age or 7 weeks for a full 12-month duration of immunity.

GROUP 2

All rabbits previously vaccinated within the last 12 months with an inactivated RHDV-2 vaccine, in addition to or without previous vaccination with Nobivac Myxo-RHD



Boost with Nobivac Myxo-RHD PLUS when next vaccine booster falls due, then boost annually.

GROUP 3

Rabbits previously vaccinated only with Nobivac Myxo-RHD (including those that have lapsed (>12 months) with vaccination for an inactivated RHDV-2 vaccine)



Vaccinate with an inactivated RHDV-2 vaccine as soon as possible.

Consider the optional use of Nobivac Myxo-RHD this year if the myxomatosis booster is also due at around the same time as the RHDV-2 vaccination.

For boosters due later this year and all subsequent boosters, use Nobivac Myxo-RHD PLUS.

Annual boosters for all groups of rabbits in all subsequent years can be managed successfully with Nobivac Myxo-RHD PLUS only.

More information is available at: www.nobivacmyxorhdplus.co.uk

Nobivac Myxo-RHD contains live myxoma vectored RHD strain 009. **POM-V**

Nobivac[®] Myxo-RHD PLUS contains live myxoma vectored RHD virus strains 009 and MK1899 for active immunisation of rabbits from 5 weeks of age onwards to reduce mortality and clinical signs of myxomatosis and rabbit haemorrhagic disease (RHD) caused by classic RHD virus (RHDV-1) and RHD type 2 virus (RHDV-2). **POM-V.**

Further information is available from the SPC, Datasheet or package leaflet.

MSD Animal Health UK Limited. Registered office Walton Manor, Walton, Milton Keynes MK7 7AJ, UK. Registered in England & Wales no. 946942.

Advice should be sought from the medicine prescriber.
Use medicines responsibly.

© 2021 MSD Animal Health UK Limited
UK-NOV-210600003 June 2021