

MIXING AND ADMINISTRATION GUIDE

GETTING STARTED

- ▶ Check the water system to ensure it works properly and is free of leaks; also ensure that water is available to all nipple or bell drinkers.
- ▶ If using a dosing pump, make sure the dosing pump is properly set to deliver the medicated water during the predetermined treatment period (hours/day).
- ▶ Determine the preferred drinking period (can range from 4 to 24 hours) when Exzolt® will be administered on the treatment day.
- ▶ Estimate how much water birds will consume during treatment based on their water consumption during the same period the day before.



CALCULATE THE DOSAGE

- ▶ **The dose is 0.5 mg fluralaner per kg bodyweight (equivalent to 0.05 ml Exzolt) administered twice, 7 days apart.**
The full course of therapy must be administered for full therapeutic effect.
- ▶ To ensure administration of the correct dosage, use calibrated scales to determine the average bodyweight within the flock.
- ▶ The required amount of product on each treatment day is calculated from the total estimated bodyweight (kg) of the entire group of chickens to be treated:
Volume of product (ml) per treatment day = Total bodyweight (kg) of chickens to be treated x 0.05 ml/kg
500 ml of Exzolt treats 10,000 kg bodyweight of birds (i.e. 5,000 chickens weighing 2 kg bodyweight) **per day** of treatment administration.

DOSING GUIDE

AVERAGE BODYWEIGHT (KG)	DAILY QUANTITY OF EXZOLT (ML) FOR THE FOLLOWING NUMBER OF BIRDS				
	1,000	5,000	10,000	15,000	20,000
1	50	250	500	750	1,000
1.1	55	275	550	825	1,100
1.2	60	300	600	900	1,200
1.3	65	325	650	975	1,300
1.4	70	350	700	1,050	1,400
1.5	75	375	750	1,125	1,500
1.6	80	400	800	1,200	1,600
1.7	85	425	850	1,275	1,700
1.8	90	450	900	1,350	1,800
1.9	95	475	950	1,425	1,900
2	100	500	1,000	1,500	2,000
2.1	105	525	1,050	1,575	2,100
2.2	110	550	1,100	1,650	2,200
2.3	115	575	1,150	1,725	2,300
2.4	120	600	1,200	1,800	2,400
2.5	125	625	1,250	1,875	2,500
3	150	750	1,500	2,250	3,000
3.5	175	875	1,750	2,625	3,500
4	200	1,000	2,000	3,000	4,000

Individual bird average bodyweight x number of birds in group x 0.05 = ml dose rate for 1 administration.

Repeat the dose 7 days later.

1 FULL TREATMENT = 2 SEPARATE DOSES, 7 DAYS APART.

PREPARING THE SOLUTION

- ▶ Mix the required volume of Exzolt with water into a large medication tank or create a stock solution diluted with drinking water and administered over time using a proportioner or dosing pump.
- ▶ When preparing the stock solution always add product and water simultaneously in order to avoid foaming.
- ▶ Rinse the container used to measure the required product volume during the filling phase ensuring that the complete dose is emptied into the medication tank or the stock solution and that no residues remain in the measuring device.
- ▶ Stir the content of the medication tank gently.
- ▶ For each day of treatment, medicated water must be freshly prepared.

START MEDICATION

- ▶ Recheck the water system to ensure it is still working properly and free of leaks; also ensure that water is available to all nipple or bell drinkers.
- ▶ Make sure the dosing pump is properly set to deliver the Exzolt during the predetermined treatment period (hours/day).
- ▶ Prime the drinker lines with medicated water and check to see when medicated water has reached the end of the line. This procedure will have to be repeated on each of the two treatment administration days.
- ▶ Check and record water consumption on each day of treatment administration and ensure all the Exzolt is consumed.

CLEAN

- ▶ After each daily treatment, resume normal water supply.



PRECAUTIONS

- ▶ Avoid contact with skin, eye and mucous membranes.
- ▶ Wear protective gloves at all times while handling the veterinary medicinal product and cleaning the measuring device. Wash hands after use. In case of skin and/or eye contact, immediately rinse with plenty of water.

Remove contaminated clothes after spillage.



Exzolt[®]
FLURALANER

Use medicines responsibly. For more information visit www.noah.co.uk/responsible

Legal category: [POM-V] Exzolt[®] is only available via your veterinary surgeon from whom advice should be sought. Exzolt[®] contains fluralaner. © Registered trademark. Exzolt[®] is the property of Intervet International B.V. or affiliated companies or licensors and is protected by copyrights, trademark and other intellectual property laws. Copyright © 2017 Intervet International B.V. All rights reserved.

Further information including side effects, precautions, warnings and contraindications is available on the product SPC or datasheet or from MSD Animal Health, Walton Manor, Walton, Milton Keynes MK7 7AJ • Tel: 0370 060 3380 • vet-support.uk@merck.com • www.msd-animal-health.co.uk

