



Nobivac® KC only protocol

Nobivac® Vaccine	6-8 WEEKS	+ 4 WEEKS	1 ST BOOSTER	YEAR 2	YEAR 3	YEAR 4	YEAR 5	
DHP			*					DHP
L4								L4
KC		†						KC

* Recommendation in line with Key Opinion Guidelines (WSAVA) & widely used in practice.¹

† Nobivac® KC can be used from 3 weeks of age.

1. Day, M. J., Horzinek, M. C., Schultz, R. D. & R. A. Squires. (2015). Guidelines for the vaccination of dogs & cats. WSAVA Vaccination Guidelines Group. JSAP 57:E1-45

Nobivac® DHP is a live attenuated vaccine containing canine distemper virus, canine adenovirus-2 and canine parvovirus. **POM-V.**

Nobivac® KC is an intranasal vaccine containing live attenuated strains of *Bordetella bronchiseptica* and canine parainfluenza virus. **POM-V.**

Nobivac® L4 is an inactivated, bacterial vaccine containing the following inactivated *Leptospira* strains: *L. interrogans* serogroup Canicola serovar Portland-vere (strain Ca-12-000), *L. interrogans* serogroup Icterohaemorrhagiae serovar Copenhageni (strain Ic-02-001), *L. interrogans* serogroup Australis serovar Bratislava (Strain As-05-073) and *L. kirschneri* serogroup Grippityphosa serovar Dadas (strain Gr-01-005). **POM-V.**

Nobivac
Protection unites us.

Further information is available from the SPC, Datasheet or package leaflet.

MSD Animal Health UK Limited. Registered office Walton Manor, Walton, Milton Keynes MK7 7AJ, UK. Registered in England & Wales no. 946942. Advice should be sought from the medicine prescriber.

Use Medicines Responsibly

© 2021 MSD Animal Health UK Ltd

UK-NOV-210200002/02/2021

MSD
Animal Health