



Nobivac® KC only protocol

Nobivac® Vaccine	6-8 WEEKS	+ 2-4 WEEKS	1 ST BOOSTER	YEAR 2	YEAR 3	YEAR 4	YEAR 5	
DHP								DHP
Lepto 2								Lepto 2
KC								KC

* Recommendation in line with Key Opinion Guidelines (WSAVA) & widely used in practice.¹

† Nobivac® KC can be used from 3 weeks of age.

^ Nobivac® DHP and Nobivac® Lepto 2 second dose from a minimum of 10 weeks of age

1. Day, M. J., Horzinek, M. C., Schultz, R. D. & R. A. Squires. (2015). Guidelines for the vaccination of dogs & cats. WSAVA Vaccination Guidelines Group. JSAP 57:E1-45

Nobivac® DHP is a live attenuated vaccine containing canine distemper virus, canine adenovirus-2 and canine parvovirus. **POM-V.**

Nobivac® KC is an intranasal vaccine containing live attenuated strains of *Bordetella bronchiseptica* and canine parainfluenza virus. **POM-V.**

Nobivac® Lepto 2 is an inactivated, bacterial vaccine containing the following inactivated *Leptospira* strains: *L. interrogans* serogroup Canicola (strain Ca-12-000), *L. interrogans* serogroup Icterohaemorrhagiae (strain 820k). **POM-V.**

Nobivac 
Protection unites us.

Further information is available from the SPC, Datasheet or package leaflet.

MSD Animal Health UK Limited. Registered office Walton Manor, Walton, Milton Keynes MK7 7AJ, UK. Registered in England & Wales no. 946942. Advice should be sought from the medicine prescriber.

Use Medicines Responsibly

© 2021 MSD Animal Health UK Ltd

UK-NOV-210200003/02/2021