



Nobivac® KC primary course followed by Nobivac® Respira Bb

Nobivac® Vaccine	6-8 WEEKS	+2-4 WEEKS	1 ST BOOSTER	YEAR 2	YEAR 3	YEAR 4	YEAR 5
DHP / DHPPi†							
Lepto 2							
Pi							
Respira Bb or KC**							

- Respira Bb
- DHPPi
- KC
- DHP
- Lepto 2
- Pi

*Recommendation in line with Key Opinion Guidelines (WSAVA) & widely used in practice.¹

† When using Nobivac® Respira Bb DHPPi and Pi are recommended to be given concurrently. Minimum finish of 10 weeks for 2nd injection of primary DHP/DHPPi course.

**Nobivac® KC can be used from 3 weeks of age.

1. Day, M. J., Horzinek, M. C., Schultz, R. D. & R. A. Squires. (2015). Guidelines for the vaccination of dogs & cats. WSAVA Vaccination Guidelines Group. JSAP 57:E1-45

Nobivac® DHP is a live attenuated vaccine containing canine distemper virus, canine adenovirus-2 and canine parvovirus. **POM-V.**

Nobivac® DHPPi is a live attenuated vaccine containing canine distemper virus, canine adenovirus-2, canine parvovirus and canine parainfluenza virus. **POM-V.**

Nobivac® KC is an intranasal vaccine containing live attenuated strains of *Bordetella bronchiseptica* and canine parainfluenza virus. **POM-V.**

Nobivac® Lepto 2 is an inactivated, bacterial vaccine containing the following inactivated *Leptospira* strains: *L. interrogans* serogroup Canicola (strain Ca-12-000), *L. interrogans* serogroup Icterohaemorrhagiae (strain 820k). **POM-V.**

Nobivac® Pi is a live attenuated vaccine containing canine parainfluenza virus. **POM-V.**

Nobivac® Respira Bb is an inactivated subunit vaccine containing the fimbrial antigen of *Bordetella bronchiseptica*. **POM-V.**

Nobivac®
 Protection unites us.

Further information is available from the SPC, Datasheet or package leaflet.

MSD Animal Health UK Limited. Registered office Walton Manor, Walton, Milton Keynes MK7 7AJ, UK. Registered in England & Wales no. 946942. Advice should be sought from the medicine prescriber.

Use Medicines Responsibly

© 2021 MSD Animal Health UK Ltd

UK-NOV-210200003/02/2021